



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Potential impact of helicobacter pylori infection on reflux disease sequence: illuminating the gap

Kountouras, Jannis ; Papaefthymiou, Apostolis ; Polyzos, Stergios A ; Liatsos, Christos ; Zavos, Christos ; Vagdatli, Eleni ; Tsikopoulos, George ; Vardaka, Elisabeth ; Tzitiridou, Maria ; Katsinelos, Panagiotis ; Doulberis, Michael

DOI: <https://doi.org/10.1097/MCG.0000000000001302>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-187567>

Journal Article

Published Version

Originally published at:

Kountouras, Jannis; Papaefthymiou, Apostolis; Polyzos, Stergios A; Liatsos, Christos; Zavos, Christos; Vagdatli, Eleni; Tsikopoulos, George; Vardaka, Elisabeth; Tzitiridou, Maria; Katsinelos, Panagiotis; Doulberis, Michael (2020). Potential impact of helicobacter pylori infection on reflux disease sequence: illuminating the gap. *Journal of Clinical Gastroenterology*, 54(2):200-201.

DOI: <https://doi.org/10.1097/MCG.0000000000001302>

Potential Impact of *Helicobacter pylori* Infection on Reflux Disease Sequence *Illuminating the Gap*

To the Editor:

Sonnenberg et al¹ reported striking variations in the ethnic distributions of gastroesophageal reflux disease (GERD) and Barrett esophagus (BE). Both diseases were associated with a lower prevalence of *Helicobacter pylori* infection (*Hp-I*) than the control population, though its prevalence in GERD and BE was higher in male than female patients, thereby running contrary to a simple consideration of an inverse relationship between *Hp-I* and these diseases.¹

In this regard, we recently suggested that epidemiological studies regarding *Hp-I*, GERD and BE-related esophageal adenocarcinoma (EAC) underestimated potential co-factors of EAC development.² Furthermore, large-scale studies concluded against the protective role of *Hp* to GERD.² The current worldwide *Hp-I* prevalence is about 58% (varied from 39.9% to 84.2%)³; its current prevalence in Asia is 66.6%. In contrast, the current global prevalence of GERD varies from 2.5% to 51.2%; its range in Asia is 33% compared with 28% of the western countries.³ This means that the conventional claim that declining *Hp* prevalence has led to a rise in GERD and its complications BE and EAC^{3,4} needs to be carefully studied. For instance, a large-scale study (~21,000 cases) reported that the decline in *Hp-I* parallels the reduction in peptic ulcer prevalence, and that the rise in GERD and/or reappearance of GERD following *Hp* therapy is rare²; contrary to expectations, patients hospitalized with duodenal ulcers (~61,500 cases), obviously attributed to *Hp-I*, had a 70% increased risk of EAC⁵; Malaysians, who for a long time have had a low prevalence of *Hp-I*, also show a low incidence of GERD, BE and distal esophageal cancers, signifying that *Hp-I* is not protective against these pathologies and its absence may be beneficial⁶; in a predominantly Caucasian

population with a high prevalence of *Hp* gastritis, *Hp-I* was not inversely connected with BE (neither presence of erosive esophagitis, length of BE nor dysplasia was associated with the presence of *Hp-I*)⁷; 2 additional studies showed that *Hp* eradication leads to better control of GERD symptoms and improves esophagitis.^{8,9} Moreover, other authors, previous supporters of the hypothesis that *Hp* “protects” against GERD, relented, claiming that *Hp* therapy does not cause or protect against GERD, and recommending *Hp* eradication in GERD.¹⁰ Such data further potentiate the consideration that *Hp* is not “protective” against anything, including GERD¹¹ and its complications BE and EAC.

Likewise, the hypothesis that increased acidity, after *Hp-I* eradication, could trigger GERD-BE-EAC sequence warrants careful reconsideration. Contrariwise, transient lower esophageal sphincter (LES) relaxations consist the substantial component of esophageal injury³ and *Hp-I* disables normal LES, probably by nitric oxide intermittent overproduction and nonselective relaxation of smooth muscles. In addition, *Hp-I* influences the gastrointestinal microbiota composition including the presence of gastric species such as *Campylobacter*; *Hp-I* induced atrophy impairs the chemical defense of stomach, thus promoting the gastric microbiota dysbiosis which might contribute to BE-EAC sequence as etc implied by the high concentrations of *Campylobacter* species in BE biofilm and the predominance of gram-negative bacteria.³ Moreover, *Hp* could directly hurt the esophagus and induce prostaglandins (PGs) overproduction that relax LES.² At molecular level, *Hp-I* induces, oncogenic gastrin and other molecular alterations which contribute to BE malignant progression.^{2,3} Specifically, *Hp-I* induced oncogenic gastrin, which, concerning its essential role in oncogenic progression in BE, stimulates proliferation via Janus Kinase (JAK) 2 and Akt-dependent nuclear factor-kappa B (NF-κB) activation in Barrett's EAC cells, exhibits an anti-apoptotic effect via Bcl-2 protein and survivin upregulations, and provokes the mitogenic and carcinogenic cyclooxygenase (COX)-2 expression. Moreover, *Hp* activates the mentioned NF-κB, a transcription regulator of inflammatory genes, including COX-2 that regulates gastrointestinal malignant cell growth and proliferation. PGs originated from upregulated COX-2 are involved in BE malignant progression, by perpetuating

chronic inflammation and the mitogenic and anti-apoptotic actions of PGs are mediated via activation of many aforementioned signaling pathways including NF-κB, Src, JAK2/STAT3, ERK, MAPK and PI3K/Akt kinases. In addition, *Hp-I* could trigger specific molecular alterations (genetic instability, E-cadherin methylation, monoclonal antibody Das-1) connected with BE pathophysiology, and stimulated Ki-67 expression predicting neoplastic progression in BE.³ Furthermore, *Hp*-related metabolic syndrome appears to contribute to the pathophysiology of GERD-BE-EAC sequence,^{12,13} and thus further research is needed to clarify this topic.

A final dilemma arises: Which cancer do we prefer to deal with, gastric cancer or EAC? The plethora of agents contributing to EAC does not allow to simplify its pathogenesis and thus the *Hp-I* positive association with gastric cancer, duodenal cancer, and colorectal cancer^{12,14} but negative association with EAC appears inconsistent, thereby warranting further investigation.

Jannis Kountouras, MD, PhD*
Apostolis Papaefthymiou, MD*†
Sergios A. Polyzos, MD, PhD*†
Christos Liatsos, MD, PhD*†
Christos Zavos, MD, PhD*
Eleni Vagdatli, MD, PhD*
George Tsikopoulos, MD, PhD*
Elisabeth Vardaka, PhD*
Maria Tzitiroidou, MD, PhD*
Panagiotis Katsinelos, MD, PhD*
Michael Doulberis, MD, DVM, PhD*§

*Department of Medicine, Second Medical Clinic, Ippokraton Hospital, Aristotle University of Thessaloniki

†Department of Pharmacology, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Macedonia

‡Department of Gastroenterology, 401 General Military Hospital of Athens Athens, Greece

§Department of Gastroenterology and Hepatology, University of Zurich, Zurich Switzerland

REFERENCES

1. Sonnenberg A, Turner KO, Genta RM. Interaction between ethnicity and *Helicobacter pylori* infection in the occurrence of reflux disease. *J Clin Gastroenterol*. 2019. [Epub ahead of print].
2. Polyzos SA, Zeglinas C, Artemaki F, et al. *Helicobacter pylori* infection and esophageal adenocarcinoma: a review and a personal view. *Ann Gastroenterol*. 2018;31:8–13.
3. Kountouras J, Doulberis M, Papaefthymiou A, et al. A perspective on risk factors

The authors declare that they have nothing to disclose.

DOI: 10.1097/MCG.0000000000001302

- for esophageal adenocarcinoma: emphasis on *Helicobacter pylori* infection. *Ann N Y Acad Sci.* 2019;1452:12–17.
4. McColl KEL. What is causing the rising incidence of esophageal adenocarcinoma in the West and will it also happen in the East? *J Gastroenterol.* 2019;54:669–673.
 5. Bahmanyar S, Zendehdel K, Nyrén O, et al. Risk of oesophageal cancer by histology among patients hospitalised for gastro-duodenal ulcers. *Gut.* 2007;56:464–468.
 6. Lee YY, Mahendra Raj S, Graham DY. *Helicobacter pylori* infection—a boon or a bane: lessons from studies in a low-prevalence population. *Helicobacter.* 2013;18:338–346.
 7. Aghayeva S, Mara KC, Katzka DA. The impact of *Helicobacter pylori* on the presence of Barrett's esophagus in Azerbaijan, a high-prevalence area of infection. *Dis Esophagus.* 2019. [Epub ahead of print].
 8. Schwizer W, Thumshirn M, Dent J, et al. *Helicobacter pylori* and symptomatic relapse of gastro-oesophageal reflux disease: a randomised controlled trial. *Lancet.* 2001;357:1738–1742.
 9. Kountouras J, Zavos C, Chatzopoulos D, et al. *Helicobacter pylori* and gastro-oesophageal reflux disease. *Lancet.* 2006;368:986; author reply 986–987.
 10. Moayyedi P. Should we test for *Helicobacter pylori* before treating gastro-oesophageal reflux disease? *Can J Gastroenterol.* 2005;19:425–427.
 11. Graham DY. *Helicobacter pylori* is not and never was “protective” against anything, including GERD. *Dig Dis Sci.* 2003;48:629–630.
 12. Kountouras J, Polyzos SA, Doulberis M, et al. Potential impact of *Helicobacter pylori*-related metabolic syndrome on upper and lower gastrointestinal tract oncogenesis. *Metabolism.* 2018;87:18–24.
 13. Doulberis M, Kountouras J, Polyzos SA, et al. Impact of *Helicobacter pylori* and/or *Helicobacter pylori*-related metabolic syndrome on gastro-oesophageal reflux disease-Barrett's esophagus-esophageal adenocarcinoma sequence. *Helicobacter.* 2018;23:e12534.
 14. Kountouras J, Doulberis M, Papaefthymiou A, et al. *Helicobacter pylori* infection and gastrointestinal tract cancer biology: considering a double-edged sword reflection. *Cell Mol Life Sci.* 2019;76:2487–2488.